

Willingness to pay report: References

- Broadbent E Petrie KJ Main J Weinman J . The Brief Illness Perception Questionnaire . J Psychosom Res 2006 ; 60 : 631 – 637 .
- Burton, N., Burton, M., Rigby, D. et al. Best-worst scaling improves measurement of first impressions. Cogn. Research 4, 36 (2019). <https://doi.org/10.1186/s41235-019-0183-2>;
- Chrzan and Patterson (2006) Chrzan, Keith and Michael Patterson (2006), "Testing for the Optimal Number of Attributes in MaxDiff Questions," 2006 Sawtooth Software Conference Proceedings, Sequim, WA.
- Cohen, Steve (2003), "Maximum Difference Scaling: Improved Measures of Importance and Preference for Segmentation," 2003 Sawtooth Software Conference Proceedings, Sequim, WA.
- Cohen, Steve and Bryan Orme (2004), "What's Your Preference?" Marketing Research, 16 (Summer 2004), 32-37.
- Devlin NJ, Shah KK, Feng Y, Mulhern B, van Hout B. Valuing health-related quality of life: An EQ-5D-5L value set for England. Health Econ. 2018 Jan;27(1):7-22. doi: 10.1002/hec.3564. Epub 2017 Aug 22. PMID: 28833869; PMCID: PMC6680214.
- Dorn, S.D., Hernandez, L., Minayas, C.B. et al. (2010). The development and validation of a new coeliac disease quality of life survey (CD-QOL). Alimentary Pharmacology and Therapeutics, 31, 666-675.
- Dubé C, Rostom A, Sy R, et al. (2005). The prevalence of celiac disease in average-risk and at-risk Western European populations: a systematic review. Gastroenterology, 128(4 Suppl 1),S57-67
- DunnGalvin, A., Barnett, J., Bejen, F.M., Ryan, K., Lucas J. (2018). Development and preliminary validation of the food intolerance quality of life questionnaire (FIQLQ): adult form. Quality of Life Research, 27, 1109-1116.
- DunnGalvin, A., Cullinane, C.D., Flokstra-de Blok, B.M.J., Dubois, A.E.J., Hourihane, J.O'B. (2010). Longitudinal validity and responsiveness of the Food Allergy Quality of Life Questionnaire – Parent Form (FAQLQ-PF) in children 0-12 years following positive and negative food challenges. Clinical and Experimental Allergy, 40, 476-85.
- DunnGalvin, A., de Blok, B.M.J., Dubois, A., Hourihane, J.O'B. (2008). Development and validation of the Food Allergy Quality of Life – Parent Administered Questionnaire (FAQLQ-PF) for food allergic children aged 0-12 years. Clinical and Experimental Allergy, 38, 977-986.
- [EQ-5D-5L | About](#)
- Feng, Y., Devlin, N. & Herdman, M. Assessing the health of the general population in England: how do the three- and five-level versions of EQ-5D compare?. Health Qual Life Outcomes 13, 171 (2015). <https://doi.org/10.1186/s12955-015-0356-8>
- Flokstra de-Blok, B.M., DunnGalvin, A., Vlieg-Boerstra, B.J., et al. (2009). Development and validation of the self-administered Food Allergy Quality of Life Questionnaire for adolescents. Journal of Allergy and Clinical Immunology, 122, 139-44.
- Hernández Alava M, Wailoo AJ, Ara R. Tails from the peak district: adjusted limited dependent variable mixture models of EQ-5D questionnaire health state utility values. Value Health. 2012 May;15(3):550-61. doi: 10.1016/j.jval.2011.12.014.
- Hernandez-Alava, M., Pudney, S., Wailoo, A. (2018). Quality review of a proposed EQ-5D-5L value set for England. Policy Research Unit in Economic Evaluation of Health and Care Interventions. Universities of Sheffield and York. EEPRU Research Report 060.
- Jansson, SA., Heibert-Arnind, M., Middelveld, R.J. et al (2013). Health-related quality of life, assessed with a disease-specific questionnaire, in Swedish adults suffering from well-

diagnosed food allergy to staple foods. *Clin Transl Allergy* 3, 21.

<https://doi.org/10.1186/2045-7022-3-21>

- Ludvigsson JF, Bai JC, Biagi F, et al. (2014). Diagnosis and management of adult coeliac disease: guidelines from the British Society of Gastroenterology. *Gut*, 63, 1210-1228
- Lins, MT.C., Tassitano, R.M., Brandt, K.G., et al (2015). Translation, cultural adaptation and validation of the celiac disease DUX (CDDUX). *Jornal de Pediatria*, 9, 448-454.
- Louviere, J. J. (1993), "The Best-Worst or Maximum Difference Measurement Model: Applications to Behavioral Research in Marketing," The American Marketing Association's 1993 Behavioral Research Conference, Phoenix, Arizona.
- [NICE Position statement on use of the EQ-5D-5L value set for England](#) (updated October 2019)
- Paez MA, Gramelspacher AM, Sinacore J et al (2017). Delay in Diagnosis of Celiac Disease in Patients Without Gastrointestinal Complaints. *Am J Med*, 130, 1318-1323
- [Sawtooth Software \(2020\) TECHNICAL PAPER SERIES The MaxDiff System Technical Paper](#) Version 9.
- Singh P, Arora A, Strand TA, Leffler DA, Catassi C, Green PH, Kelly CP, Ahuja V, Makharia GK. (2018). Global Prevalence of Celiac Disease: Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol*. 2018.
- Szende A, Janssen B, Cabases J (2014). [Self-Reported Population Health: An International Perspective based on EQ-5D](#). Springer.
- Train, K. (2009). *Discrete Choice Methods with Simulation*. Cambridge: Cambridge University Press.
- Van Doorn, R.K., Winkler, L.M.F., Zwinderman, K.H. et al (2008). CDDUX: A Disease-specific Health-related Quality-of-life Questionnaire for Children with Celiac Disease. *Journal of Pediatric Gastroenterology and Nutrition*, 47, 147-152.
- Zipser RD, Patel S, Yahya KZ et al. (2003). Presentations of adult celiac disease in a nationwide patient support group. *Dig Dis Sci*, 48(4),761-4.