

# Risk from *Listeria monocytogenes* in ready to eat smoked fish: References

- Aalto-Araneda, M. et al. (2019) 'Processing plant and machinery sanitation and hygiene practices associate with *Listeria monocytogenes* occurrence in ready-to-eat fish products', *Food Microbiology*, 82, pp. 455–464. doi:10.1016/j.fm.2019.03.017.
- Abdelhamed, H. et al. (2019) 'Validation of Predicted Virulence Factors in *Listeria monocytogenes* Identified Using Comparative Genomics', *Toxins*, 11(9), p. 508. doi:10.3390/toxins11090508.
- Acciari, V.A. et al. (2017) '*Listeria monocytogenes* in Smoked Salmon and Other Smoked Fish at Retail in Italy: Frequency of Contamination and Strain Characterization in Products from Different Manufacturers', *Journal of Food Protection*, 80(2), pp. 271–278. doi:10.4315/0362-028X.JFP-15-599.
- ACMSF (2020) [Fixed-term group on multidimensional representation of risks \(PDF\)](#).
- Adak, G.K., Long, S.M. and O'Brien, S.J. (2002) 'Trends in indigenous foodborne disease and deaths, England and Wales: 1992 to 2000', *Gut*, 51(6), pp. 832–841. doi:10.1136/gut.51.6.832.
- Aureli, P. et al. (2000) 'An outbreak of febrile gastroenteritis associated with corn contaminated by *Listeria monocytogenes*', *The New England Journal of Medicine*, 342(17), pp. 1236–1241. doi:10.1056/NEJM200004273421702.
- Bader, G., Al-Tarawneh, M. and Myers, J. (2016) 'Review of Prosthetic Joint Infection from *Listeria monocytogenes*', *Surgical Infections*, 17(6), pp. 739–744. doi:10.1089/sur.2016.067.
- Branciarri, R. et al. (2016) 'Thermal inactivation and growth potential of *Listeria monocytogenes* in smoked tench', *Italian Journal of Food Safety*, 5(3). doi:10.4081/ijfs.2016.5974.
- Brennan, M. et al. (2013) [Activity Recognition and Temperature Monitoring \(ART\) Feasibility Study. FS244026 \(PDF\)](#).
- Chasseignaux, E. et al. (2001) 'Molecular epidemiology of *Listeria monocytogenes* isolates collected from the environment, raw meat and raw products in two poultry- and pork-processing plants', *Journal of Applied Microbiology*, 91(5), pp. 888–899. doi:10.1046/j.1365-2672.2001.01445.x.
- Chau, M.L. et al. (2017) 'Microbial survey of ready-to-eat salad ingredients sold at retail reveals the occurrence and the persistence of *Listeria monocytogenes* Sequence Types 2 and 87 in pre-packed smoked salmon', *BMC Microbiology*, 17(1), p. 46. doi:10.1186/s12866-017-0956-z.
- [Chilled Seafood in Multiple Retail \(2021\) Seafish \(PDF\)](#) (Accessed: 25 May 2022).
- [ComBase \(no date\)](#). (Accessed: 17 May 2022).
- Dalton, C.B. et al. (1997) 'An Outbreak of Gastroenteritis and Fever Due to *Listeria monocytogenes* in Milk', *New England Journal of Medicine*, 336(2), pp. 100–106. doi:10.1056/NEJM199701093360204.
- Dominguez, C., Gomez, I. and Zumalacarregui, J. (2001) 'Prevalence and Contamination Levels of *Listeria monocytogenes* in Smoked Fish and Pâté Sold in Spain', *Journal of Food Protection*, 64(12), pp. 2075–2077. doi:10.4315/0362-028X-64.12.2075.
- Drevets, D.A. and Bronze, M.S. (2008) '*Listeria monocytogenes*: epidemiology, human disease, and mechanisms of brain invasion', *FEMS immunology and medical microbiology*, 53(2), pp. 151–165. doi:10.1111/j.1574-695X.2008.00404.x.
- EFSA (2013) 'Analysis of the baseline survey on the prevalence of *Listeria monocytogenes* in certain ready-to-eat foods in the EU, 2010-2011 [Part A: \*Listeria monocytogenes\*](#)

- [prevalence estimates](#), *EFSA Journal*, 11(6). (Accessed: 5 May 2022).
- EFSA (2014) 'Analysis of the baseline survey on the prevalence of *Listeria monocytogenes* in certain ready-to-eat foods in the EU, 2010–2011 [Part B: analysis of factors related to prevalence and exploring compliance](#)', *EFSA Journal*, 12(8).
  - EFSA (2018) '[Listeria monocytogenes contamination of ready-to-eat foods and the risk for human health in the EU](#)', *EFSA Journal*, 16(1).
  - EFSA (2020) [United Kingdom- Trends and Sources of Zoonoses and Zoonotic Agents in Foodstuffs, Animals and Feedingstuffs in 2020. \(PDF\)](#)
  - EFSA and ECDC (2018) 'Multi-country outbreak of *Listeria monocytogenes* sequence type 8 infections linked to consumption of salmon products', *EFSA Supporting Publications*, 15(10), p. 1496E. doi:10.2903/sp.efsa.2018.EN-1496.
  - EFSA and ECDC (2019) 'Multi-country outbreak of *Listeria monocytogenes* clonal complex 8 infections linked to consumption of cold-smoked fish products', *EFSA Supporting Publications*, 16(6), p. 1665E. doi:10.2903/sp.efsa.2019.EN-1665.
  - Eicher, C. et al. (2020) 'Growth Potential of *Listeria monocytogenes* in Three Different Salmon Products', p. 13.
  - Eklund, M.W. et al. (1995) 'Incidence and Sources of *Listeria monocytogenes* in Cold-Smoked Fishery Products and Processing Plants', *Journal of Food Protection*, 58(5), pp. 502–508. doi:10.4315/0362-028X-58.5.502.
  - Ericsson, H. (1997) 'An Outbreak of Listeriosis Suspected To Have Been Caused by Rainbow Trout', 35, p. 4.
  - European Salmon Smokers Association (2018) '[European Guide to Good Practice for Smoked and/or Salted and/or Marinated Fish](#)'. (Accessed: 6 May 2022).
  - [Everything You Need to Know About ... Smoked Mackerel - Fresh Fish Daily \(no date\)](#). (Accessed: 25 May 2022).
  - FAO (1997) [Hazard Analysis and Critical Control Point \(HACCP\) System and Guidelines for its Application](#). (Accessed: 6 May 2022).
  - FAO (2006) [The Use of Microbiological Risk Assessment Outputs to Develop Practical Risk Management Strategies: Metrics to improve food safety: Report](#). Rome, Italy: FAO. (Accessed: 6 May 2022).
  - FAO and WHO (2006) '[Development of Practical Risk Management Strategies based on Microbiological Risk Assessment Outputs](#)'.(PDF)
  - Farber, J.M. et al. (2021) 'Alternative approaches to the risk management of *Listeria monocytogenes* in low risk foods', *Food Control*, 123, p. 107601. doi:10.1016/j.foodcont.2020.107601.
  - Farber, J.M. and Peterkin, P.I. (1991) '*Listeria monocytogenes*, a food-borne pathogen', *Microbiological Reviews*, 55(3), pp. 476–511. doi:10.1128/mr.55.3.476-511.1991.
  - FDA (2003) '[Quantitative Assessment of Relative Risk to Public Health from Foodborne \*Listeria monocytogenes\* Among Selected Categories of Ready-to-Eat Foods](#)', FDA [Preprint].(Accessed: 21 April 2022).
  - Food Standards Australia New Zealand (2016) '[Imported food risk statement: Processed ready-to-eat finfish and \*Listeria monocytogenes\*](#)'. (PDF).
  - FSA (2008) '[A microbiological survey of retail smoked fish with particular reference to the presence of \*Listeria monocytogenes\*](#)'.
  - FSA (2018) '[Reducing the risk of vulnerable groups contracting Listeriosis: Guidance for healthcare and social care organisations](#)'. (PDF).
  - FSA (2021) '[Incident Management Plan for Non-Routine Incidents](#)'. (PDF) (Accessed: 15 May 2022).
  - FSA (2022) 'The risk to vulnerable consumers from *Listeria monocytogenes* in blue cheese'.
  - FSS (2018) [Reducing the risk of vulnerable groups contracting listeriosis. \(PDF\)](#) Accessed: 30 May 2022).
  - FSS (no date) [Safe Smoked Fish Tool](#). (Accessed: 17 May 2022).
  - Fuentes, A. et al. (2010) 'Physicochemical Characterization of Some Smoked and Marinated Fish Products', *Journal of Food Processing and Preservation*, 34(1), pp. 83–103.

doi:10.1111/j.1745-4549.2008.00350.x.

- Gahan, C.G.M. and Hill, C. (2005) 'Gastrointestinal phase of *Listeria monocytogenes* infection', *Journal of Applied Microbiology*, 98(6), pp. 1345–1353. doi:10.1111/j.1365-2672.2005.02559.x.
- Gillesberg Lassen, S. et al. (2016) 'Two listeria outbreaks caused by smoked fish consumption—using whole-genome sequencing for outbreak investigations', *Clinical Microbiology and Infection*, 22(7), pp. 620–624. doi:10.1016/j.cmi.2016.04.017.
- González-Rodríguez, M.-N. et al. (2002) 'Numbers and types of microorganisms in vacuum-packed cold-smoked freshwater fish at the retail level', *International Journal of Food Microbiology*, 77(1–2), pp. 161–168. doi:10.1016/s0168-1605(02)00048-x.
- Guillet, C. et al. (2010) 'Human Listeriosis Caused by *Listeria ivanovii*', *Emerging Infectious Diseases*, 16(1), pp. 136–138. doi:10.3201/eid1601.091155.
- [Health Protection Agency \(2009\) Guidelines for Assessing the Microbiological Safety of Ready-to-Eat Foods Placed on the Market, Guidelines for Assessing the Microbiological Safety of Ready-to-Eat Foods Placed on the Market. \(PDF\)](#)
- ['Hot smoked, cold smoked - what's the difference?' \(2019\)](#) Mere Trout, 24 November. (Accessed: 25 May 2022).
- ['Hot smoked mackerel \(no date\)](#). (Accessed: 25 May 2022).
- Jami, M. et al. (2014) '*Listeria monocytogenes* in Aquatic Food Products—A Review', *Comprehensive Reviews in Food Science and Food Safety*, 13(5), pp. 798–813. doi:10.1111/1541-4337.12092.
- Jemmi, T. and Keusch, A. (1992) 'Behavior of *Listeria monocytogenes* during processing and storage of experimentally contaminated hot-smoked trout', *International Journal of Food Microbiology*, 15(3), pp. 339–346. doi:10.1016/0168-1605(92)90067-D.
- Kang, J. et al. (2012) 'Effect of curing method and freeze-thawing on subsequent growth of *Listeria monocytogenes* on cold-smoked salmon', *Journal of Food Protection*, 75(9), pp. 1619–1626. doi:10.4315/0362-028X.JFP-11-561.
- Lachmann, R. et al. (2022) 'Invasive listeriosis outbreaks and salmon products: a genomic, epidemiological study', *Emerging Microbes & Infections*, 0(ja), pp. 1–30. doi:10.1080/22221751.2022.2063075.
- Lambertz, S.T. et al. (2012) 'Prevalence and level of *Listeria monocytogenes* in ready-to-eat foods in Sweden 2010', *International Journal of Food Microbiology*, 160(1), pp. 24–31. doi:10.1016/j.ijfoodmicro.2012.09.010.
- Lappi, V.R. et al. (2004) 'Prevalence and growth of *Listeria* on naturally contaminated smoked salmon over 28 days of storage at 4 degrees C', *Journal of Food Protection*, 67(5), pp. 1022–1026. doi:10.4315/0362-028x-67.5.1022.
- Lindqvist, R. and Westö, A. (2000) 'Quantitative risk assessment for *Listeria monocytogenes* in smoked or gravad salmon and rainbow trout in Sweden', *International Journal of Food Microbiology*, 58(3), pp. 181–196. doi:10.1016/S0168-1605(00)00272-5.
- Liu, D. et al. (2005) 'Comparative assessment of acid, alkali and salt tolerance in *Listeria monocytogenes* virulent and avirulent strains', *FEMS Microbiology Letters*, 243(2), pp. 373–378. doi:10.1016/j.femsle.2004.12.025.
- Lopez-Valladares, G., Danielsson-Tham, M.-L. and Tham, W. (2018) 'Implicated Food Products for Listeriosis and Changes in Serovars of *Listeria monocytogenes* Affecting Humans in Recent Decades', *Foodborne Pathogens and Disease*, 15(7), pp. 387–397. doi:10.1089/fpd.2017.2419.
- Lyhs, U. et al. (2001) 'Microbiological quality and shelf-life of vacuum-packaged "gravad" rainbow trout stored at 3 and 8 degrees C', *International Journal of Food Microbiology*, 70(3), pp. 221–230. doi:10.1016/s0168-1605(01)00548-7.
- Mead, P.S. et al. (1999) 'Food-related illness and death in the United States', *Emerging Infectious Diseases*, 5(5), pp. 607–625. doi:10.3201/eid0505.990502.
- Miettinen, M.K. et al. (1999) 'Molecular Epidemiology of an Outbreak of Febrile Gastroenteritis Caused by *Listeria monocytogenes* in Cold-Smoked Rainbow Trout', *Journal of Clinical Microbiology*, 37(7), pp. 2358–2360. doi:10.1128/JCM.37.7.2358-2360.1999.

- Miller, A.J. (1992) 'Combined Water Activity and Solute Effects on Growth and Survival of *Listeria monocytogenes* Scott A', *Journal of Food Protection*, 55(6), pp. 414–418. doi:10.4315/0362-028X-55.6.414.
- Mowi (2021) '[Salmon Farming Industry Handbook](#)'. (PDF).
- Nakari, U.-M. et al. (2014) 'Investigation of increased listeriosis revealed two fishery production plants with persistent *Listeria* contamination in Finland in 2010', *Epidemiology & Infection*, 142(11), pp. 2261–2269. doi:10.1017/S095026881300349X.
- [National Diet and Nutrition Survey \(no date\) Food Standards Agency](#). (Accessed: 15 May 2022).
- National Fisheries Institute and National Food Processors Association (2002) '[Smoked Seafood Working Group: \*Listeria monocytogenes\* Control Manual](#)'. (PDF)
- Neunlist, M.R. et al. (2005) 'Effect of salting and cold-smoking process on the culturability, viability, and virulence of *Listeria monocytogenes* strain Scott A', *Journal of Food Protection*, 68(1), pp. 85–91. doi:10.4315/0362-028x-68.1.85.
- [NHS \(2020\) Foods to avoid in pregnancy, nhs.uk](#). (Accessed: 30 May 2022).
- [NHS 111 Wales \(2022\) Pregnancy Guide- Foods to Avoid](#). (Accessed: 30 May 2022).
- [NHS Inform \(2022\) Eating well in pregnancy](#). (Accessed: 30 May 2022).
- [NIDirect \(2022\) Foods to avoid in pregnancy](#). (Accessed: 30 May 2022).
- [Office for National Statistics \(no date\) Population estimates for the UK, England and Wales, Scotland and Northern Ireland: mid-2020](#). (Accessed: 17 May 2022).
- Orsi, R.H. and Wiedmann, M. (2016) 'Characteristics and distribution of *Listeria* spp., including *Listeria* species newly described since 2009', *Applied Microbiology and Biotechnology*, 100, pp. 5273–5287. doi:10.1007/s00253-016-7552-2.
- Peck, M. et al. (2006) [Clostridium botulinum in vacuum packed \(VP\) and modified atmosphere packed \(MAP\) chilled foods](#). B13006.
- Pezdirc, K.B. et al. (2012) '*Listeria monocytogenes* and diet during pregnancy; balancing nutrient intake adequacy v. adverse pregnancy outcomes', *Public Health Nutrition*, 15(12), pp. 2202–2209. doi:10.1017/S1368980012000717.
- PHE (2017) '*Listeriosis* in England and Wales: Summary for 2017', p. 16.
- [PHE \(2018\) Listeriosis in England and Wales: summary for 2018, GOV.UK](#). (Accessed: 20 January 2022).
- PHE (2021) [Listeriosis in England and Wales: summary for 2019, GOV.UK](#). (Accessed: 24 May 2022).
- PHS (2020) [Annual summary of Listeria infections, 2019](#). (Accessed: 22 April 2022).
- Porsby, C.H. et al. (2008) 'Influence of processing steps in cold-smoked salmon production on survival and growth of persistent and presumed non-persistent *Listeria monocytogenes*', *International Journal of Food Microbiology*, 122(3), pp. 287–295. doi:10.1016/j.ijfoodmicro.2008.01.010.
- Pouillot, R. et al. (2009) 'Quantitative Risk Assessment of *Listeria monocytogenes* in French Cold-Smoked Salmon: II. Risk Characterization', *Risk Analysis*, 29(6), pp. 806–819. doi:10.1111/j.1539-6924.2008.01200.x.
- Pouillot, R. et al. (2015) '*Listeria monocytogenes* Dose Response Revisited—Incorporating Adjustments for Variability in Strain Virulence and Host Susceptibility', *Risk Analysis*, 35(1), pp. 90–108. doi:10.1111/risa.12235.
- Pouillot, R. et al. (2016) 'Infectious Dose of *Listeria monocytogenes* in Outbreak Linked to Ice Cream, United States, 2015', *Emerging Infectious Diseases*, 22(12). doi:10.3201/eid2212.160165.
- Poysky, F.T. et al. (1997) 'Inactivation of *Listeria monocytogenes* on Hot-smoked Salmon by the Interaction of Heat and Smoke or Liquid Smoke †', *Journal of Food Protection*, 60(6), pp. 649–654. doi:10.4315/0362-028X-60.6.649.
- Rasmussen, R.R. et al. (2017) 'Effects of industrial processing on essential elements and regulated and emerging contaminant levels in seafood', *Food and Chemical Toxicology*, 104, pp. 85–94. doi:10.1016/j.fct.2017.02.008.
- Ricci, A. et al. (2017) 'Public health risks associated with hepatitis E virus (HEV) as a food-borne pathogen', *Efsa Journal*, 15(7), p. 4886. doi:10.2903/j.efsa.2017.4886.



- Rørvik, L.M. (2000) 'Listeria monocytogenes in the smoked salmon industry', International Journal of Food Microbiology, p. 8.
- Rørvik, L.M., Yndestad, M. and Skjerve, E. (1991) 'Growth of Listeria monocytogenes in vacuum-packed, smoked salmon, during storage at 4° C', International Journal of Food Microbiology, 14(2), pp. 111–117. doi:10.1016/0168-1605(91)90097-9.
- Rotariu, O. et al. (2014) 'Smoked salmon industry practices and their association with Listeria monocytogenes', Food Control, 35(1), pp. 284–292. doi:10.1016/j.foodcont.2013.07.015.
- Sandercock, H. (2019) [What lies beneath: fish category report 2019, The Grocer](#). (Accessed: 30 May 2022).
- Schjørring, S. et al. (2017) 'Cross-border outbreak of listeriosis caused by cold-smoked salmon, revealed by integrated surveillance and whole genome sequencing (WGS), Denmark and France, 2015 to 2017', p. 5.
- Tham, W. et al. (2000) 'Lessons from an outbreak of listeriosis related to vacuum-packed gravad and cold-smoked fish', International Journal of Food Microbiology, 62(3), pp. 173–175. doi:10.1016/S0168-1605(00)00332-9.
- Thomas, D.J.I. et al. (2012) [A review of the published literature and current production and processing practices in smoked fish processing plants with emphasis on contamination by Listeria monocytogenesA. FS425012. \(PDF\)](#)
- Tocmo, R. et al. (2014) 'Listeria monocytogenes in Vacuum-Packed Smoked Fish Products: Occurrence, Routes of Contamination, and Potential Intervention Measures', Comprehensive Reviews in Food Science and Food Safety, 13(2), pp. 172–189. doi:10.1111/1541-4337.12052.
- Townsend, A. et al. (2021) 'A Systematic Review of Listeria Species and Listeria monocytogenes Prevalence, Persistence, and Diversity throughout the Fresh Produce Supply Chain', Foods, 10(6), p. 1427. doi:10.3390/foods10061427.
- UK Health Security Agency (no date) [Listeria: enhanced surveillance questionnaire, GOV.UK](#). (Accessed: 17 May 2022).
- Uyttendaele, M. et al. (2009) 'Prevalence and challenge tests of Listeria monocytogenes in Belgian produced and retailed mayonnaise-based deli-salads, cooked meat products and smoked fish between 2005 and 2007', International Journal of Food Microbiology, 133(1–2), pp. 94–104. doi:10.1016/j.ijfoodmicro.2009.05.002.
- Van Coillie, E. et al. (2004) 'Prevalence and Typing of Listeria monocytogenes in Ready-to-Eat Food Products on the Belgian Market', Journal of Food Protection, 67(11), pp. 2480–2487. doi:10.4315/0362-028X-67.11.2480.
- Walland, J. et al. (2015) 'Listeria monocytogenes infection in ruminants: Is there a link to the environment, food and human health? A review', Schweizer Archiv Fur Tierheilkunde, 157(6), pp. 319–328. doi:10.17236/sat00022.
- Warriner, K. and Namvar, A. (2009) 'What is the hysteria with Listeria?', Trends in Food Science & Technology, 20(6), pp. 245–254. doi:10.1016/j.tifs.2009.03.008.
- White, K. (2019) [Fresh: Meat, Fish & Poultry: Top Products 2019, The Grocer](#). (Accessed: 30 May 2022).
- WHO (2004) [Risk assessment of Listeria monocytogenes in ready-to-eat foods: technical report](#).
- [WHO \(2018\) Listeriosis](#), (Accessed: 22 April 2022).
- Zhelyazkov, G. and Stratev, D. (2018) 'Some Physicochemical Characteristics of Fish Products Sampled from Bulgarian Retail Markets', Journal of food quality and hazards control, 5(1), pp. 33–36.