

# Risk of campylobacteriosis from low-throughput poultry slaughterhouses:

## References

- ACMSF (2019) [Third report on Campylobacter](#). (Accessed: 16 September 2022).
- Allen, V.M. et al. (2007) 'Campylobacter spp. contamination of chicken carcasses during processing in relation to flock colonisation', *International Journal of Food Microbiology*, 113(1), pp. 54–61. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2006.07.011>.
- Allen, V.M. et al. (2008) 'Sources and Spread of Thermophilic Campylobacter spp. during Partial Depopulation of Broiler Chicken Flocks', *Journal of Food Protection*, 71(2), pp. 264–270. Available at: <https://doi.org/10.4315/0362-028X-71.2.264>.
- Ansari-Lari, M. et al. (2011) 'Prevalence and risk factors associated with campylobacter infections in broiler flocks in Shiraz, southern Iran', *International Journal of Food Microbiology*, 144(3), pp. 475–479. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2010.11.003>.
- Antic, D. (2022) 'Personal communication'.
- Ausvet Europe et al. (2022) Modelling framework to quantify the risk of AMR exposure via food products - example of chicken and lettuce. Food Standards Agency. Available at: <https://doi.org/10.46756/sci.fsa.gum110>.
- Baranyi, J. and Tamplin, M.L. (2004) 'ComBase: A Common Database on Microbial Responses to Food Environments', *Journal of Food Protection*, 67(9), pp. 1967–1971. Available at: <https://doi.org/10.4315/0362-028X-67.9.1967>.
- Battersby, T., Whyte, P. and Bolton, D. (2016) 'Protecting broilers against Campylobacter infection by preventing direct contact between farm staff and broilers', *Food Control*, 69, pp. 346–351. Available at: <https://doi.org/10.1016/j.foodcont.2016.04.053>.
- Biglia, A. et al. (2018) 'Temperature and energy performance of domestic cold appliances in households in England', *International Journal of Refrigeration*, 87, pp. 172–184. Available at: <https://doi.org/10.1016/j.ijrefrig.2017.10.022>.
- Borck Høg, B. et al. (2016) 'Farm specific risk factors for Campylobacter colonisation in Danish and Norwegian broilers', *Preventive Veterinary Medicine*, 130, pp. 137–145. Available at: <https://doi.org/10.1016/j.prevetmed.2016.04.002>.
- Boysen, L., Knøchel, S. and Rosenquist, H. (2007) 'Survival of Campylobacter jejuni in different gas mixtures', *FEMS Microbiology Letters*, 266(2), pp. 152–157. Available at: <https://doi.org/10.1111/j.1574-6968.2006.00525.x>.
- Bruhn, C.M. (2014) 'Chicken Preparation in the Home: An Observational Study', *Food Protection Trends*, 34(5), pp. 318–330.
- Chowdhury, S. et al. (2012) 'Risk factors for Campylobacter infection in Danish broiler chickens', *Poultry Science*, 91(10), pp. 2701–2709. Available at: <https://doi.org/10.3382/ps.2012-02412>.
- Collineau, L. et al. (2020) 'A farm-to-fork quantitative risk assessment model for Salmonella Heidelberg resistant to third-generation cephalosporins in broiler chickens in Canada', *International Journal of Food Microbiology*, 330, p. 108559. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2020.108559>.
- Corry, J. et al. (2017) 'Reducing campylobacter cross-contamination during poultry processing', p. 159.
- Corry, J.E.L. et al. (2007) 'Surface pasteurisation of chicken carcasses using hot water', *Journal of Food Engineering*, 79(3), pp. 913–919. Available at: <https://doi.org/10.1016/j.jfoodeng.2006.03.018>.

- Davis, L. and DiRita, V. (2008) 'Growth and laboratory maintenance of *Campylobacter jejuni*', *Current Protocols in Microbiology*, Chapter 8, p. Unit 8A.1.1-8A.1.7. Available at: <https://doi.org/10.1002/9780471729259.mc08a01s10>.
- Didier, P. et al. (2021) 'Washing hands and risk of cross-contamination during chicken preparation among domestic practitioners in five European countries', *Food Control*, 127, p. 108062. Available at: <https://doi.org/10.1016/j.foodcont.2021.108062>.
- Dogan, O.B. et al. (2019) 'A quantitative microbial risk assessment model of *Campylobacter* in broiler chickens: Evaluating processing interventions', *Food Control*, 100, pp. 97–110. Available at: <https://doi.org/10.1016/j.foodcont.2019.01.003>.
- EFSA Panel on Biological Hazards (BIOHAZ) (2011) 'Scientific Opinion on *Campylobacter* in broiler meat production: control options and performance objectives and/or targets at different stages of the food chain', *EFSA Journal*, 9(4), p. 2105. Available at: <https://doi.org/10.2903/j.efsa.2011.2105>.
- EFSA Panel on Biological Hazards (BIOHAZ) et al. (2020) 'Update and review of control options for *Campylobacter* in broilers at primary production', *EFSA journal*. European Food Safety Authority, 18(4), p. e06090. Available at: <https://doi.org/10.2903/j.efsa.2020.6090>.
- Ellis-Iversen, J. et al. (2009) 'Risk factors for *Campylobacter* colonisation during rearing of broiler flocks in Great Britain', *Preventive Veterinary Medicine*, 89(3–4), pp. 178–184. Available at: <https://doi.org/10.1016/j.prevetmed.2009.02.004>.
- EVANS, E.W. and REDMOND, E.C. (2016) 'Time-Temperature Profiling of United Kingdom Consumers' Domestic Refrigerators', *Journal of Food Protection*, 79(12), pp. 2119–2127. Available at: <https://doi.org/10.4315/0362-028X.JFP-16-270>.
- FSA (2019a) [Campylobacter Reduction Programme: Update, Food Standards Agency](#). (Accessed: 17 November 2022).
- FSA (2019b) Meat Industry Guide - Chapter 13.
- FSA (2022) 'Personal communication'.
- Georgiev, M., Beauvais, W. and Guitian, J. (2017) 'Effect of enhanced biosecurity and selected on-farm factors on *Campylobacter* colonization of chicken broilers', *Epidemiology and Infection*, 145(3), pp. 553–567. Available at: <https://doi.org/10.1017/S095026881600251X>.
- Guirin, G.F. et al. (2020) 'Prevalence of *Campylobacter jejuni* and *Campylobacter coli* from broilers at conventional and kosher abattoirs and retail stores', *Revista Argentina de Microbiología*, 52(3), pp. 217–220. Available at: <https://doi.org/10.1016/j.ram.2019.07.002>.
- Hastings, R. et al. (2011) 'Campylobacter genotypes from poultry transportation crates indicate a source of contamination and transmission', *Journal of Applied Microbiology*, 110(1), pp. 266–276. Available at: <https://doi.org/10.1111/j.1365-2672.2010.04883.x>.
- Holland, D. et al. (2020) 'Estimating deaths from foodborne disease in the UK for 11 key pathogens', *BMJ open gastroenterology*, 7(1), p. e000377. Available at: <https://doi.org/10.1136/bmjgast-2020-000377>.
- Holland, D. and Mahmoudzadeh, N. (2020) Foodborne Disease Estimates for the United Kingdom in 2018. Food Standards Agency. Available at: <https://doi.org/10.46756/sci.fsa.squ824>.
- Huneau-Salaün, A. et al. (2007) 'Risk factors for *Campylobacter* spp. colonization in French free-range broiler-chicken flocks at the end of the indoor rearing period', *Preventive Veterinary Medicine*, 80(1), pp. 34–48. Available at: <https://doi.org/10.1016/j.prevetmed.2007.02.001>.
- Hutchison, M. et al. (2016) [Monitoring of campylobacters in UK poultry slaughter batches and carcasses and the collection of information from primary production and processing for risk factor elucidation](#).(PDF)
- IPSOS Mori (2016) [Trialling a modernised poultry inspection system in UK slaughterhouses](#)
- James, C. et al. (2007) 'Decontamination of poultry carcasses using steam or hot water in combination with rapid cooling, chilling or freezing of carcass surfaces', *International Journal of Food Microbiology*, 114(2), pp. 195–203. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2006.09.019>.

- Jorgensen, F. et al. (2011) 'Influence of Season and Geography on Campylobacter jejuni and C. coli Subtypes in Housed Broiler Flocks Reared in Great Britain', *Applied and Environmental Microbiology*, 77(11), pp. 3741–3748. Available at: <https://doi.org/10.1128/AEM.02444-10>.
- Jorgensen, F. et al. (2019) 'A microbiological survey of Campylobacter contamination in fresh whole UK-produced chilled chickens at retail sale', p. 37. <https://doi.org/10.46756/sci.fsa.xx2973>
- Kaakoush, N.O. et al. (2015) 'Global Epidemiology of Campylobacter Infection', *Clinical Microbiology Reviews*, 28(3), pp. 687–720. Available at: <https://doi.org/10.1128/CMR.00006-15>.
- Kaakoush, N.O., Mitchell, H.M. and Man, S.M. (2015) 'Chapter 67 - Campylobacter', in Y.-W. Tang et al. (eds) *Molecular Medical Microbiology (Second Edition)*. Boston: Academic Press, pp. 1187–1236. Available at: <https://doi.org/10.1016/B978-0-12-397169-2.00067-6>.
- McDowell, S.W.J. et al. (2008) 'Campylobacter spp. in conventional broiler flocks in Northern Ireland: epidemiology and risk factors', *Preventive Veterinary Medicine*, 84(3–4), pp. 261–276. Available at: <https://doi.org/10.1016/j.prevetmed.2007.12.010>.
- Nastasijevic, I. et al. (2020) 'The European Union control strategy for Campylobacter spp. in the broiler meat chain', *Journal of Food Safety*, 40(5), p. e12819. Available at: <https://doi.org/10.1111/jfs.12819>.
- Näther, G. et al. (2009) 'Analysis of risk factors for Campylobacter species infection in broiler flocks', *Poultry Science*, 88(6), pp. 1299–1305. Available at: <https://doi.org/10.3382/ps.2008-00389>.
- O'Brien, S.J. et al. (2016) 'Modelling study to estimate the health burden of foodborne diseases: cases, general practice consultations and hospitalisations in the UK, 2009', *BMJ Open*, 6(9), p. e011119. Available at: <https://doi.org/10.1136/bmjopen-2016-011119>.
- Oxford University (2021) 'Enhanced molecular-based surveillance and source attribution of campylobacter infections in the UK', Food Standards Agency [Preprint]. Available at: <https://doi.org/10.46756/sci.fsa.ksj135>.
- PHE (2021) 'A microbiological survey of campylobacter contamination in fresh whole UK-produced chilled chickens at retail sale (Y6)', Food Standards Agency [Preprint]. Available at: <https://doi.org/10.46756/sci.fsa.xx2973>.
- Rasschaert, G. et al. (2020) 'Campylobacter contamination of broilers: the role of transport and slaughterhouse', *International Journal of Food Microbiology*, 322, p. 108564. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2020.108564>.
- Refrégier-Petton, J. et al. (2001) 'Risk factors for Campylobacter spp. contamination in French broiler-chicken flocks at the end of the rearing period', *Preventive Veterinary Medicine*, 50(1–2), pp. 89–100. Available at: [https://doi.org/10.1016/s0167-5877\(01\)00220-3](https://doi.org/10.1016/s0167-5877(01)00220-3).
- Robinson, D.A. (1981) 'Infective dose of Campylobacter jejuni in milk.', *BMJ*, 282(6276), pp. 1584–1584. Available at: <https://doi.org/10.1136/bmj.282.6276.1584>.
- Rosenquist, H. et al. (2006) 'The effect of slaughter operations on the contamination of chicken carcasses with thermotolerant Campylobacter', *International Journal of Food Microbiology*, 108(2), pp. 226–232. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2005.12.007>.
- Tam, C.C. et al. (2012) 'Longitudinal study of infectious intestinal disease in the UK (IID2 study): incidence in the community and presenting to general practice', *Gut*, 61(1), pp. 69–77. Available at: <https://doi.org/10.1136/gut.2011.238386>.
- Teunis, P.F.M. et al. (2018) 'Acute illness from Campylobacter jejuni may require high doses while infection occurs at low doses', *Epidemics*, 24, pp. 1–20. Available at: <https://doi.org/10.1016/j.epidem.2018.02.001>.
- Torralbo, A. et al. (2014) 'Prevalence and risk factors of Campylobacter infection in broiler flocks from southern Spain', *Preventive Veterinary Medicine*, 114(2), pp. 106–113. Available at: <https://doi.org/10.1016/j.prevetmed.2014.01.019>.
- Tustin, J. et al. (2011) 'A national epidemic of campylobacteriosis in Iceland, lessons learned', *Zoonoses and Public Health*, 58(6), pp. 440–447. Available at:

<https://doi.org/10.1111/j.1863-2378.2010.01387.x>.

- V. Allain et al. (2014) 'Prevalence of and risk factors for Campylobacter colonisation in broiler flocks at the end of the rearing period in France', *British poultry science*, 55(4), pp. 452–459. Available at: <https://doi.org/10.1080/00071668.2014.941788>.
- Van Limbergen, T. et al. (2018) 'Scoring biosecurity in European conventional broiler production', *POULTRY SCIENCE*, 97(1), pp. 74–83. Available at: <https://doi.org/10.3382/ps/pex296>.
- Vose Consulting (US) LLC (2011) [A quantitative microbiological risk assessment of Campylobacter in the broiler meat chain](#) | EFSA. (Accessed: 6 October 2022).
- Weis, A.M. et al. (2016) 'Genomic Comparison of Campylobacter spp. and Their Potential for Zoonotic Transmission between Birds, Primates, and Livestock', *Applied and Environmental Microbiology*, 82(24), pp. 7165–7175. Available at: <https://doi.org/10.1128/AEM.01746-16>.